

Prognosis of IntraCerebral Haemorrhage (PITCH)

Principal Investigator

Charlotte Cordonnier

Centre Hospitalier Régional Universitaire de Lille,
Department of Neurology & Stroke Unit,
Hopital Roger Salengro,
59037 Lille, France.

Charlotte.cordonnier@chru-lille.fr

Summary

Country	France
Principal Investigator	Charlotte Cordonnier
Contact person (email)	Charlotte.cordonnier@chru-lille.fr <
Key publication/reference	Cordonnier et al. Brain 2010; 133: 3281-9
Years in which study conducted	2004-2009
Sample	
Size	562
Population: Hospital/community	Hospital
Selection: consecutive/random	Consecutive
Admit with previous stroke?	Yes
Admit with TIA?	No
Age range	Median 72 (IQR 58-79); no age limit for inclusion
Number of centres	1
Control group: number, population, selection	None
Assessment	
Initial: Time and data collected/tests administered	At admission: MedHx, VRF Day 7: Neurological exam, glucose, renal, hb, white cells, cholesterol, ECG, CT, MRI, IQCODE

Detailed	Discharge: function
Subsequent (follow-ups)	NΨ: 6m, 1y, 2y, 3y, 4.5y, 6y, 8y, 10y
Stroke-related data	NIHSS
Functional tests/data	Barthel Index, Rankin Scale
Other medical tests/data	Fatigue, seizures
Neuropsychological tests	Test battery, MMSE, Mattis
MRI scans, when and how many	At admission, 6m, 2y, 4.5y, 6y, 8y, 10y
PET scans	No
Psychiatric exams/diagnoses	Dementia, anxiety, depression (MADRS)
Intervention trialled?	No

CT=computed tomography scan, MedHx=medical history, VRF=vascular risk factors (hypertension, diabetes, atrial fibrillation, obesity, smoking etc.), NΨ=neuropsychological, TIA=transient ischemic attack, m=month, y=year

